

<i>Product Licence Number</i>	<i>Company Name</i>	<i>Product Name</i>	<i>Active Ingredients</i>	<i>Indications</i>	<i>Date of Authorisation</i>
0011/0195 (continued)	John Wyeth and Brother Limited T/A Wyeth Laboratories			3. Therapeutic supplementation in conditions such as osteomalacia, rickets, post-gastrectomy malabsorption, or dietary deficiencies. 4. Where dietary requirement is high as in childhood, pregnancy and lactation due to increase demand, or in old age due to impaired absorption.	
0012/0222	May & Baker Limited T/A Rhone-Poulenc Rorer	Toprek Tablets 25 mg	Ketoprofen BP 25 mg	Prescription Only Medicine Symptomatic treatment of mild to moderate pain (such as dental, post-operative and postpartum pain, dysmenorrhoea and headache.	28th May 1993
0012/0259	May & Baker Limited T/A Rhone-Poulenc Rorer	Zolerim 7.5 mg Tablets	Zopiclone HSE 7.5 mg	Prescription Only Medicine <i>Recommended Clinical Indications</i> For the short term treatment of insomnia, including difficulties in falling asleep, nocturnal awakening and early awakening, transient, situational or chronic insomnia and insomnia secondary to psychiatric disturbances, in situations where the insomnia is debilitating or is causing severe distress for the patient. Long term continuous use is not recommended. A course of treatment should employ the lowest effective dose and should be no longer than 4 weeks in duration.	6th May 1993
0022/0134	Kabi Pharmacia Limited	Dipentum Capsules	Olsalazine Sodium 250 mg	Prescription Only Medicine Oral treatment of mild acute ulcerative colitis and maintenance of remission.	13th May 1993
0032/0180	Upjohn Limited	Colestid Tablets 1g	Colestipol Hydrochloride USP 1 gm	Prescription Only Medicine Colestid is indicated as adjunctive therapy to diet in the management of patients with elevated cholesterol levels who have not responded adequately to diet. It may be used alone or in combination with additional lipid lowering agents. Colestid may elevate serum triglyceride levels when used as sole therapy. This elevation is generally transient but may persist in some individuals. Although Colestid is effective in all types of hypercholesterolaemia, it is medically most appropriate in patients with Fredrickson's type II hyperlipoproteinaemia. Dietary therapy specific for the type of hypercholesterolaemia should be the initial treatment of choice. Excess body weight may be an important factor and weight reduction should be attempted prior to drug therapy in the over-weight. The use of drugs should be considered only when reasonable attempts have been made to obtain satisfactory results with non-drug methods. When drug therapy is begun, the patient should be instructed of the importance of adhering to the correct diet.	24th May 1993
0037/0234	Abbott Laboratories Limited	Hytrin BPH 1 mg Tablets	Terazosin Monohydrochloride HSE 1.19 mg	Prescription Only Medicine As a therapy for the symptomatic treatment of urinary obstruction caused by benign prostatic hyperplasia (BPH).	20th May 1993
0037/0235	Abbott Laboratories Limited	Hytrin BPH 2 mg Tablets	Terazosin Monohydrochloride HSE 2.37 mg	Prescription Only Medicine Same as in PL 0037/0234	20th May 1993
0037/0236	Abbott Laboratories Limited	Hytrin BPH 5 mg Tablets	Terazosin Monohydrochloride HSE 5.94 mg	Prescription Only Medicine Same as in PL 0037/0234	20th May 1993
0037/0237	Abbott Laboratories Limited	Hytrin BPH 10 mg Tablets	Terazosin Monohydrochloride HSE 11.87 mg	Prescription Only Medicine Same as in PL 0037/0234	20th May 1993
0071/0405	Sterling-Winthrop Group Limited T/A Sterling Health	Andrews Seltzer	Paracetamol EP 500.0 mg Sodium Bicarbonate EP 1342.0 mg	General Sale List - may be sold or supplied otherwise than by or under the supervision of a pharmacist only when in packs of not more than 25 tablets. Andrews Seltzer is particularly effective for headache and upset stomach, dyspepsia or acid indigestion and will also help relieve the pain and discomfort of period pain, migraine, rheumatic pain, toothache, colds, flu and a sore throat.	4th May 1993