

Product Licence Number	Company Name	Product Name	Active Ingredients	Indications	Date of Authorisation
12724/0076	GM Laboratories Limited	Diazepam Tablets BP 2 mg	Diazepam BP 2.00 mg	<p>2. Abnormal Retention of Fluids Acetazolamide acts on the reversible hydration of carbon dioxide and hydration of the carbonic acid reaction in the kidney resulting in renal loss of bicarbonate ion which carries out sodium, water and potassium. It can be used in conjunction with other diuretics when effects on several segments of the nephron are desirable in the treatment of fluid retaining states.</p> <p>3. Epilepsy When used in conjunction with other anticonvulsants best results have been seen in petit mal in children. Good results have been seen in both children and adults with grand mal, mixed seizure patterns and myoclonic jerk patterns.</p>	1st November 1994
12724/0077	GM Laboratories Limited	Diazepam Tablets BP 5mg	Diazepam BP 5.00 mg	<p>Prescription Only Medicine</p> <p>1. Relief of Anxiety For the short-term relief (2-4 weeks) only of anxiety that is severe, disabling or subjecting the individual to unacceptable distress, occurring alone or in association with insomnia or short-term psychosomatic, organic or psychotic illness. The use of benzodiazepines to treat short-term 'mild' anxiety is considered to be inappropriate and unsuitable.</p> <p>2. Insomnia To treat insomnia only when it is severe, disabling, or subjecting the individual to extreme distress.</p> <p>3. Premedication</p> <p>4. Control of Muscle Spasm.</p> <p>Prescription Only Medicine</p> <p>1. Relief of Anxiety Benzodiazepines are indicated for the short-term relief (2-4 weeks) only of anxiety that is severe, disabling or subjecting the individual to unacceptable distress, occurring alone or in association with insomnia or short-term psychosomatic, organic psychotic illness. The use of benzodiazepines to treat short-term 'mild' anxiety is considered to be inappropriate and unsuitable.</p> <p>2. Insomnia Benzodiazepines should be used to treat insomnia only when it is severe, disabling, or subjecting the individual to extreme distress.</p> <p>3. Premedication</p> <p>4. Control of Muscle Spasm.</p>	1st November