

Product Licence Number	Company Name	Product Name	Active Ingredients	Indications	Date of Authorisation
				ankylosing spondylitis, acute musculoskeletal disorders, degenerative joint disease of the hip, low back pain and acute gouty arthritis. It is also indicated in inflammation, pain and oedema following orthopaedic procedures; and the treatment of pain and associated symptoms of primary dysmenorrhoea. Indocid suppositories may be used where night pain and morning stiffness are prominent. One suppository at bedtime will frequently give relief from pain and stiffness for 13 to 16 hours after administration.	
01502/0054	Phoenix Pharmaceuticals Limited	Sterile Dobutamine Hydrochloride Injection Concentrate 250mg/20ml	Dobutamine Hydrochloride USP 1.4% w/v equivalent to Dobutamine 1.25% w/v	Prescription Only Medicine It is indicated in adults who require inotropic support in the treatment of low output cardiac failure associated with myocardial infarction, open heart surgery, cardiomyopathies, septic shock and cardiogenic shock. It can also increase or maintain cardiac output during positive end expiratory pressure (PEEP) ventilation. It may also be used for cardiac stress testing as an alternative exercise in patients for whom routine exercise cannot be satisfactorily performed.	16th December 1994
1664/0001	Apothekernes Laboratorium AS	Link Suspension 15%	Aluminium Hydroxide-Magnesium Carbonate co-dried gel HSE 750mg	General Sale List For the treatment of (a) Dyspepsia, heartburn, gastric discomfort, (b) Symptomatic treatment of hyperacidity, gastritis, oesophagitis, hiatus hernia. To promote healing of gastric and duodenal ulcers.	1st October 1994
1664/0002	Apothekernes Laboratorium AS	Link Suspension 30%	Aluminium Hydroxide-Magnesium Carbonate co-dried gel HSE 1.5 gm	General Sale List See PL/1664/0001	1st October 1994
1664/0003	Apothekernes Laboratorium AS	Apolar 0.1% Cream	Desonide 1.0 mg	Prescription Only Medicine Eczema and dermatitis of all types including atopic eczema, photodermatitis, primary irritant and allergic dermatitis, lichen planus, lichen simplex, prurigo nodularis.	1st October 1994
1664/0004	Apothekernes Laboratorium AS	Link Tablets 700 mg	Aluminium Hydroxide-Magnesium Carbonate co-dried gel HSE 700mg	General Sale List See PL/1664/0001	1st October 1994
1817/0016	Mallinckrodt Medical (UK) Limited	TechneScan® MAG3	Benzoylmercaptoacetyltiglycine (Betiatide) HSE 1 mg	Prescription Only Medicine A diagnostic agent to be used in the evaluation of nephrological and urological disorders.	7th October 1994
02848/0176	Antigen International Limited	Vancomycin Hydrochloride for Injection 500 mg	Vancomycin base equivalent to 500,000 i.u. of Vancomycin base	Prescription Only Medicine It is indicated for the treatment of potentially life-threatening infections which can not be treated with other less toxic microbial agents, including penicillins and cephalosporins. It may be used to treat severe staphylococcal infections in patients who cannot receive or who have failed to respond to penicillins and cephalosporins, or infections with staphylococci resistant to other antibiotics. It is used in the treatment of endocarditis and in the prophylaxis of endocarditis in patients at risk from dental or surgical procedures. Its effectiveness has also been demonstrated in other infections due to Staphylococci, including Osteomyelitis, Pneumonia, Septicaemia and soft tissue infections.	16th January 1995
03433/0147	Farmitalia Carlo Erba Limited	Farlutal Oral Suspension 100mg/ml	Medroxyprogesterone Acetate BP 500.0 mg	Prescription Only Medicine Palliative treatment of hormone-sensitive malignancies. It has been	6th October 1994