

DEPARTMENT OF HEALTH MEDICINES CONTROL AGENCY

Market Towers, 1 Nine Elms Lane, London SW8 5NQ. Telephone 01-720 2188, Enquiries to Ext 3375 or 3373

Product Licences Granted

<i>Product Licence Number</i>	<i>Company Name</i>	<i>Product Name</i>	<i>Active Ingredients</i>	<i>Indications (Non-Prescription Items)</i>
PL 0001/0111	Ciba-Geigy Plc	Proflex Cream	Ibuprofen BP 5.0% w/w	Proflex Cream is a topical analgesic and anti-inflammatory treatment for the fast relief of the symptoms of rheumatic and muscular pain, backache, sprains, strains, lumbago and fibrositis. <i>Route of Administration:</i> Topical POM
PL 0009/0069	Pharmacia Limited UK	Dipentum Capsules	Olsalazine Sodium 250 mg	POM
PL 0017/0238	Astra Pharmaceuticals Limited	Losec Capsules	Omeprazole 20.00 mg	POM
PL 0019/0080	William R. Warner and Company Limited	Parkol Elixir	Paracetamol PHE 120.0 mg	<i>Indications:</i> To relieve pain and fever associated with teething, earache, headache, colds and influenza and general aches and pains in children. <i>Route of Administration:</i> Oral PM The product is indicated for the treatment and prevention of athletes foot (<i>tinea pedis</i>). It may also be helpful in dhobie itch (<i>tinea cruris</i>) and prickly heat (miliaria). <i>Route of Administration</i> The foot ointment is for topical application to the skin. GSL
PL 0039/0266	Evans Medical Limited	Mycil Ointment	Tolnaftate BP 1.0% w/w, Benzalkonium Chloride Solution BP 0.2% v/w	The product is indicated for the adjunctive treatment and prevention of athletes foot (<i>tinea pedis</i>). It may also be helpful in conditions where tenderness and sweating cause skin irritation. Examples of these conditions are dhobie itch (<i>tinea cruris</i>) and prickly heat (miliaria). <i>Route of Administration</i> The dry powder spray is for topical application to the skin. GSL
PL 0039/0301	Evans Medical Limited	Mycil Athlete's Foot Spray	Tolnaftate BP 0.64% w/w	For the treatment of plantar and other viral warts with the exception of warts on the face and those in the anal and perineal region. To be applied to the wart twice daily or as directed
PL 0041/0015	Bioglan Laboratories Limited	Novaruca ^(R)	Glutaraldehyde BP 10% w/w, as 50% Solution 20% w/w	