

Product Licence Number	Company Name	Product Name	Active Ingredients	Indications	Date of Authorisation
12724/0076	GM Laboratories Limited	Diazepam Tablets BP 2 mg	Diazepam BP 2.00 mg	<p>glaucoma and perioperatively in acute angle closure glaucoma where delay of surgery is desired in order to lower intraocular pressure. Acetazolamide acts on inflow, decreasing the amount of aqueous secretion.</p> <p>2. <i>Abnormal retention of fluids</i>—Acetazolamide acts on the reversible hydration of carbon dioxide and hydration of the carbonic acid reaction in the kidney resulting in renal loss of bicarbonate ion which carries out sodium, water and potassium. It can be used in conjunction with other diuretics when effects on several segments of the nephron are desirable in the treatment of fluid retaining states.</p> <p>3. <i>Epilepsy</i>—When used in conjunction with other anticonvulsants best results have been seen in petit mal in children. Good results have been seen in both children and adults with grand mal, mixed seizure patterns and myoclonic jerk patterns.</p> <p>POM</p> <p>1. <i>Relief of anxiety</i>—for the short-term relief (2-4 weeks) only of anxiety that is severe, disabling or subjecting the individual to unacceptable distress, occurring alone or in association with insomnia or short-term psychosomatic, organic or psychotic illness. The use of benzodiazepines to treat short-term "mild" anxiety is considered to be inappropriate and unsuitable.</p> <p>2. <i>Insomnia</i>—To treat insomnia only when it is severe, disabling, or subjecting the individual to extreme distress.</p> <p>3. Premedication.</p> <p>4. <i>Control of muscle spasm.</i></p> <p>POM</p>	1st November 1994
12724/0077	GM Laboratories Limited	Diazepam Tablets BP 5mg	Diazepam BP 5.00mg	<p>1. Relief of Anxiety</p> <p>Benzodiazepines are indicated for the short-term relief (2-4 weeks) only of anxiety that is severe, disabling or subjecting the individual to unacceptable distress, occurring alone or in association with insomnia or short-term psychosomatic, organic or psychotic illness.</p> <p>The use of Benzodiazepines to treat short-term 'mild' anxiety is considered to be inappropriate and unsuitable.</p> <p>2. <i>Insomnia</i></p> <p>Benzodiazepines should be used to treat insomnia only when it is severe, disabling, or subjecting the individual to extreme distress.</p> <p>3. Premedication.</p> <p>4. Control of Muscle Spasm.</p> <p>POM</p>	1st November 1994
12724/0078	GM Laboratories Limited	Diazepam Tablets BP 10mg	Diazepam BP 10.00mg	See PL 12724/0077	1st November 1994
12724/0079	GM Laboratories Limited	Co-Trimoxazole Tablets BP	Trimethoprim BP 80.00 mg Sulphamethoxazole BP 400.00 mg	<p>Effective against a wide range of gram positive and gram negative organisms. Of value in the treatment of the following:</p> <p>Respiratory Tract: Acute and Chronic Bronchitis, Lobar and Bronchopneumonia, Otitis Media and Sinusitis.</p> <p>Genito-Urinary Tract: Urethritis, Cystitis, Pyelonephritis and Prostatitis, Male and Female Gonorrhoea.</p> <p>Gastro-Intestinal Tract: Typhoid and Paratyphoid Fevers, Chronic Carriage of <i>Salmonella</i> Typhi and Paratyphi, Cholera (as an adjunct to fluid and electrolyte replacement), Shigellosis.</p> <p>Skin Infections: Abscesses and Wound Infections.</p> <p>POM</p>	1st November 1994
12724/0080	GM Laboratories Limited	Ephedrine Hydrochloride Tablets BP 30mg	Ephedrine Hydrochloride BP 30.00mg	For the treatment of reversible airways obstruction when other remedies fail.	1st November 1994
12724/0082	GM Laboratories Limited	Calcium Gluconate Tablets BP 600mg	Calcium Gluconate BP 600.00mg	For the prophylaxis of calcium deficiency in persons who may not get enough calcium in their regular diet and also for the treatment of hypocalcaemia occurring in conditions such as chronic hypoparathyroidism,	1st November 1994